

**Evidence-Based Complementary and Alternative Medicine**



Home Editorial Board Author Guidelines Aims and Scope Submit a Manuscript

Evid Based Complement Alternat Med. 2015; 2015: 874849.

PMCID: PMC4363882

Published online 2015 Mar 2. doi: [10.1155/2015/874849](https://doi.org/10.1155/2015/874849)

## Self-Reported Efficacy of Cannabis and Other Complementary Medicine Modalities by Parkinson's Disease Patients in Colorado

Taylor Andrew Finseth,<sup>1</sup> Jessica Louise Hedeman,<sup>1</sup> Robert Preston Brown, II,<sup>2</sup> Kristina I. Johnson,<sup>3</sup> Matthew Sean Binder,<sup>1</sup> and Benzi M. Kluger<sup>1,\*</sup>

<sup>1</sup>Department of Neurology, University of Colorado Anschutz Medical Campus, Aurora, CO 80045, USA

<sup>2</sup>Department of Economics, University of Colorado Denver, Denver, CO 80217, USA

<sup>3</sup>Colorado School of Public Health, University of Northern Colorado, Greeley, CO 80639, USA

\*Benzi M. Kluger: Email: [benzi.kluger@ucdenver.edu](mailto:benzi.kluger@ucdenver.edu)

Academic Editor: Wei-Bo Zhang

Received 2014 Dec 16; Revised 2015 Feb 17; Accepted 2015 Feb 19.

Copyright © 2015 Taylor Andrew Finseth et al.

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

Go to:

**Introduction.** Complementary and alternative medicine (CAM) is frequently used by Parkinson's disease (PD) patients. We sought to provide information on CAM use and efficacy in PD patients in the Denver metro area with particular attention to cannabis use given its recent change in legal status. **Methods.** Self-administered surveys on CAM use and efficacy were completed by PD patients identified in clinics and support groups across the Denver metro area between 2012 and 2013. **Results.** 207 patients (age  $69 \pm 11$ ; 60% male) completed the survey. Responses to individual CAM therapy items showed that 85% of respondents used at least one form of CAM. The most frequently reported CAMs were vitamins (66%), prayer (59%), massage (45%), and relaxation (32%). Self-reported improvement related to the use of CAM was highest for massage, art therapy, music therapy, and cannabis. While only 4.3% of our survey responders reported use of cannabis, it ranked among the most effective CAM therapies. **Conclusions.** Overall, our cross-sectional study was notable for a high rate of CAM utilization amongst PD patients and high rates of self-reported efficacy across most CAM modalities. Cannabis was rarely used in our population but users reported high efficacy, mainly for nonmotor symptoms.

### 1. Introduction

Go to:

Complementary and alternative medicine (CAM) is defined by the National Center of Complementary and Alternative Medicine as a group of diverse medical and healthcare systems, practices, and products that are not considered an integral part of conventional medicine [1]. CAM modalities include alternative medical systems like traditional Chinese medicine; bioactive substances like vitamins or herbal supplements; manipulative therapies like massage or chiropractic; mind-body therapies like yoga or relaxation; and energy therapies like Reiki. CAM is estimated to be used by approximately 38% of the US population with an estimated 34 billion dollars spent on CAM products and services annually [2, 3].

Parkinson's disease (PD) is the second most common neurodegenerative disorder following Alzheimer's disease and leads to impairments in function and quality of life through both motor and nonmotor symptoms. A recent review article examining the use of CAM in PD found that the usage rates and most commonly used modalities by

country were as follows: South Korea (76%, oriental medicine), USA (40%, vitamins), UK (39%, massage), Sweden (34%, acupuncture), and Argentina (26%, acupuncture) [4].

The last study of CAM use by PD patients in the USA was conducted in 2001 on outpatients of a tertiary academic practice in Baltimore [5]. Our study sought to provide updated figures on the prevalence and spectrum of CAM use in PD patients in Colorado across multiple providers and to provide data on self-reported efficacy of CAM for various symptoms.

In addition, we specifically investigated usage of cannabis amongst the PD population in Colorado. Medical cannabis has been legally available with authorization from a physician in Colorado since 2000 and available to patients in retail outlets with a prescription since 2010. Recreational use of cannabis without a prescription was legalized in Colorado in November 2012, just prior to the start of our study. There have been few clinical trials of cannabis in PD, most of which were underpowered with uncontrolled series suggesting benefit for both motor and nonmotor symptoms and controlled trials with negative or inconclusive results [6]. In contrast, a single survey based study from the Czech Republic reported high rates of use (25%) and benefit for multiple symptoms (>40%) [7]. Although this came from a large sample ( $N = 339$ ) there was a 54% nonresponse rate raising issues of reporting bias. Studies of medical cannabis in the general population show frequent use for pain, insomnia, and anxiety/depression, all of which are common nonmotor symptoms of PD [8]. Due to the paucity of literature on the subject and potential interest and benefit in our PD population, we sought to gather information on cannabis usage and self-reported efficacy amongst the Colorado PD population.

## 2. Methods

Go to:

### 2.1. Standard Protocol Approvals, Registrations, and Patient Consents

The Colorado Multiple Institutional Review Board approved this study and, given that it was an anonymous survey not collecting identifying information, waived consent requirements.

### 2.2. Patient Surveys

A self-administered survey was provided to patients at the University of Colorado Hospital Movement Disorders Clinics and PD support groups in the Denver metro area between November 2012 and August 2013. The anonymous survey was completed either in person or at the patient's convenience and returned to the study coordinator in person or via mail.

The survey asked patients to indicate past or current use of individual CAM modalities provided in a list format for management of symptoms associated with Parkinson's disease (see Supplementary Material for complete survey available online at <http://dx.doi.org/10.1155/2015/874849>). If they had used a modality, they were asked to answer follow-up questions of whether they had noticed "great improvement," "some improvement," "no improvement," or "worsening/side effects" with that modality. They were also asked if the modalities used had been associated with improvement in any of the following areas: quality of life, mood, sleep, energy, or motor symptoms. The survey collected self-reported demographic information, disease characteristics. Data on multivitamins, diet, exercise, and physical therapy were also collected for comparison purpose. Exercise and physical therapy were not considered CAM as they are considered standard of care for PD [9]. Multivitamins and diet were not considered CAM for purposes of estimating CAM prevalence because they are commonly employed by the general public for general health maintenance. We did not ask about the specific use of diet for PD or use of specific diets (e.g., gluten-free).

Incomplete surveys were not discarded for purposes of analysis. Completion rates for the majority of items ranged from 95 to 97% with the exception of income, which had a 74% completion rate. Three of the 210 completed surveys were discarded due to suspected inaccuracy of demographic information: one with age listed as 2006 and two surveys with initial diagnosis at ages 18 and 26. We did not track refusal rate, but we estimate it was below

10% and typically due to time constraints.

### 2.3. Statistical Analysis

Study data were input and managed using Research Electronic Data Capture (REDCap) hosted at the University of Colorado Denver Anschutz Medical Campus [10]. Statistical analyses were performed using SAS 9.3 (Cary, NC) and included unadjusted descriptive statistics. Determination of associations between demographic and clinical variables and CAM use was performed using Student's *t*-test for continuous variables and Chi-square test or Fisher's exact test (if cell frequencies were less than 5) for categorical variables. *P* < 0.05 was considered statistically significant.

## 3. Results

Go to:

### 3.1. Demographics

Two hundred and seven patients with PD completed the survey between November 2012 and August 2013. [Table 1](#) describes the demographics of our cohort. 82% had at least some college education and 33% had graduate level education. A majority of the patients were of higher socioeconomic status with 60% reporting income above \$60,000. Average duration of Parkinson's symptoms was 8.15 ( $\pm 6.9$ ) years.

Table 1	
Characteristics of the study population	
Age, <i>n</i> (mean $\pm$ SD)	60.8 ( $\pm$ 10.6)
Male, <i>n</i> (%)	103 (49.5)
Female, <i>n</i> (%)	104 (49.5)
Race:	
White, <i>n</i> (%)	196 (94.6)
Asian/Pacific Islander, <i>n</i> (%)	1 (0.5)
African American, <i>n</i> (%)	1 (0.5)
Middle Eastern, <i>n</i> (%)	1 (0.5)
Hispanic, <i>n</i> (%)	1 (0.5)
Other, <i>n</i> (%)	1 (0.5)
Education:	
Less than high school, <i>n</i> (%)	33 (15.7)
High school, <i>n</i> (%)	66 (31.4)
Some college, <i>n</i> (%)	65 (30.8)
Graduate, <i>n</i> (%)	43 (20.1)
Income:	
Less than \$60,000, <i>n</i> (%)	62 (29.4)
\$60,000+, <i>n</i> (%)	145 (69.6)
Socioeconomic status:	
Low income, <i>n</i> (%)	103 (49.5)
Medium income, <i>n</i> (%)	104 (49.5)

[Table 1](#)

Characteristics of the study population.

### 3.2. Estimated Prevalence and Self-Reported Efficacy of CAM Use

52% of patients reported past or current use of CAM therapies for treatment of PD symptoms. However, when surveys were analyzed based on specific CAM therapy items, 85% of patients demonstrated current or recent use of at least one form of CAM ([Table 2](#)). 88% of responders reported that their physician was aware of their CAM use. Vitamins, not including multivitamins, were the most frequently used CAM (66%), of which vitamin E and CoQ10 were the most commonly used. Other common forms of CAM included prayer (49%), massage (45%), and relaxation techniques (32%).

Table 2	
Most frequently used CAM treatments <sup>a</sup>	
Rank	Treatment
1	Vitamin
2	Prayer
3	Massage
4	Acupuncture
5	Relaxation
6	Supplements
7	Meditation
8	Music
9	Yoga
10	Chiropractic

[Table 2](#)

Most frequently used CAM treatments<sup>a</sup>.

We further analyzed self-reported efficacy of various CAMs as well as several commonly accepted nonpharmacological therapies such as physical therapy and exercise ([Table 3](#)). Self-reported improvement with CAM modalities varied from 29% to 86% and was similar to well accepted non-CAM therapies. Most therapies had a response rate (defined by report of some or great improvement due to the therapy) of 60% or greater. Massage and meditation had the highest response rates of the commonly used therapies. Music therapy, cannabis, and chiropractic also had high rates of reported effectiveness, although in a smaller proportion of the total sample. The CAM therapies with the highest rate of reported improvement in specific domains are listed in [Table 3](#). Acupuncture, exercise, homeopathy, and spiritual therapy each had one respondent report worsening or side effects. Side effects were not reported with any other therapy.

Table 3	
Most efficacious CAM treatments <sup>a</sup>	
Rank	Therapy
1	Physical therapy
2	Music
3	Supplements
4	Exercise
5	Acupuncture
6	Homeopathy
7	Chiropractic
8	Prayer
9	Massage
10	Relaxation
11	Supplements
12	Medication
13	Supplements
14	Supplements
15	Supplements
16	Supplements
17	Supplements
18	Supplements
19	Supplements
20	Supplements
21	Supplements
22	Supplements
23	Supplements
24	Supplements
25	Supplements
26	Supplements
27	Supplements
28	Supplements
29	Supplements
30	Supplements
31	Supplements
32	Supplements
33	Supplements
34	Supplements
35	Supplements
36	Supplements
37	Supplements
38	Supplements
39	Supplements
40	Supplements
41	Supplements
42	Supplements
43	Supplements
44	Supplements
45	Supplements
46	Supplements
47	Supplements
48	Supplements
49	Supplements
50	Supplements
51	Supplements
52	Supplements
53	Supplements
54	Supplements
55	Supplements
56	Supplements
57	Supplements
58	Supplements
59	Supplements
60	Supplements
61	Supplements
62	Supplements
63	Supplements
64	Supplements
65	Supplements
66	Supplements
67	Supplements
68	Supplements
69	Supplements
70	Supplements
71	Supplements
72	Supplements
73	Supplements
74	Supplements
75	Supplements
76	Supplements
77	Supplements
78	Supplements
79	Supplements
80	Supplements
81	Supplements
82	Supplements
83	Supplements
84	Supplements
85	Supplements
86	Supplements
87	Supplements
88	Supplements
89	Supplements
90	Supplements
91	Supplements
92	Supplements
93	Supplements
94	Supplements
95	Supplements
96	Supplements
97	Supplements
98	Supplements
99	Supplements
100	Supplements

[Table 3](#)

Most efficacious CAM treatments<sup>a</sup>.

Nine (4.3%) responders reported use of cannabis to treat their PD symptoms. Of those nine, four had fourteen or more years of education and eight were white. Ages ranged from 49 to 75, and these patients reported PD symptom duration ranging from 2 to 11 years. Seven rated their overall health as good or excellent, while two rated their overall health as fair. Five reported “great improvement” in their symptoms with cannabis. In particular, five reported improvement in their mood and sleep, and two reported improvement in their motor symptoms and quality of life. No one reported worsening of symptoms or side effects.

### 3.3. Association of Demographic/Clinical Variables and CAM Use

There were no associations found between demographic or disease characteristics and overall CAM use. Regarding the five most common forms of CAM, females were significantly more likely to utilize massage, relaxation techniques, and acupuncture and African Americans and Asians were significantly more likely to utilize relaxation techniques. No significant correlations were found for age, education, or income with the five most commonly reported forms of CAM. Disease characteristics of years since diagnosis, age at initial diagnosis, number of PD medications used, and current health status were also examined without significant correlations with the top 5 most common forms of CAM. There were no significant differences in the demographics of patients using cannabis compared to demographics of all survey respondents.

## 4. Discussion

Go to:

Our study was the first to systematically examine self-reported efficacy of CAM therapies in PD patients including cannabis. We found higher CAM usage rates than prior studies for the general US population (28–54%) and similar or higher rates reported for other neurological disorders like stroke (30%), sciatica (47%), memory loss (47%), migraines (50%), epilepsy (70%), pediatric neurology (78%), and multiple sclerosis (81%) [11–15]. In addition, our results suggest a higher percentage of PD patients use CAM than that found by Rajendran et al., who estimated 40% CAM use amongst a similar cohort of PD patients in Maryland [5]. Due to the large discrepancy between the findings of these two studies, it is worthwhile to explore possible explanations. One explanation may be regional differences. The Western United States has been shown to have higher CAM usage rates amongst the general population compared to other regions [16, 17]. Both Denver and Boulder have a very visible complementary and alternative medicine community, which may lead to increased awareness and access to CAM. The two studies also used different techniques; Rajendran et al. interviewed consecutive patients of a single practitioner with low refusal rate. Our study, in contrast, studied patients from multiple practitioners at both academic centers and community support groups, data was self-reported anonymously, and refusal rate was higher. This design is more prone to overestimate the true prevalence of CAM use given possibility of self-selection bias. Despite this possibility, we believe the true prevalence of CAM use amongst our population is likely higher than that found by Rajendran et al. and may reflect regional variability as well as societal trends of increasing CAM use in the past decade [5]. Given this suspected regional variability, our study's findings may not accurately reflect CAM usage patterns amongst PD patients on a national level.

The relative use of the various CAM modalities in the USA is similar to that found by Rajendran et al. in 2001. Vitamins and herbal supplements remain the most utilized, followed by prayer, massage, relaxation, and acupuncture. Our findings also demonstrate differences in CAM use by PD patients in Colorado when compared to PD patients in other countries [4]. While acupuncture is the most commonly used CAM therapy for PD worldwide, it is just the fifth most popular CAM in our cohort which may be reflective of low insurance coverage of acupuncture services in the USA. In contrast, vitamin usage in the USA greatly exceeds that in any other region studied to date [18–20]. The reasons for high rates of vitamin usage in the USA may be explained by their wide availability and ease of use. We also suspect that part of the discrepancy between self-reported rates of CAM use and calculated rates may be due to patients not considering vitamins as a form of CAM. Despite their popularity, vitamins had low reported efficacy in improvement in quality of life or other metrics compared to other CAM modalities. Notably, the two most commonly used vitamins (vitamin E and CoQ10) have both been shown in large

randomized clinical trials to not improve PD outcomes [21, 22].

Self-reported efficacy for many CAM therapies was high for both motor and nonmotor symptoms. Reported improvement in certain domains suggests areas where further clinical research may be useful. Meditation, art therapy, and cannabis all had responder rates of greater than 50% for mood improvement and, with the exception of cannabis, might be considered to augment standard pharmacotherapy or psychotherapeutic approaches. In the author's experience, cannabis is potentially helpful for anxiety but not depression and there is in fact evidence for cannabis use increasing depressive symptoms [23]. Cannabis had the highest responder rate for sleep improvement at 55% and might be considered for refractory insomnia not responsive to first-line agents. Motor improvement rates approaching 50% were reported by patients using massage, chiropractic, and tai chi. Notably, there is strong evidence from randomized clinical trials to support tai chi use and consistent but less rigorous data for massage [24–26]. Chiropractic has not been studied and caution may be warranted for manipulations, particularly involving the cervical spine [27].

The use of cannabis as a therapeutic agent for various medical conditions has been well documented, but there are few studies looking specifically at its use for treating PD symptoms [6, 28–31]. While only a small number of participants in our study reported use of cannabis for PD, those that did reported benefits in mood (56%), sleep (56%), motor symptoms (22%), and quality of life (22%). Our results differ from those found in a survey of Czech PD patients from 2004 in several important ways [28]. Usage rates are lower (4.3% versus 25.1%) and motor response was lower (22% versus 44.7%). However, the few randomized clinical trials performed to date have failed to substantiate any objective benefits and suggest side effects including lightheadedness and drowsiness are fairly common [32]. Our data suggests that PD patients in our cohort tend to utilize cannabis for its impact on nonmotor symptoms rather than motor symptoms, and it was rated as the most effective therapy for sleep and mood improvement amongst all CAMs. There is a single small observational study also suggesting that cannabis therapies may be helpful for psychosis in PD patients with dementia [33]. While the role of cannabis in PD is not yet clearly delineated we suspect its use will grow in popularity given the recent legalization in select states, the wide range of potential formulations in these states, and increased media attention. Potential side effects of cognitive impairment, apathy, fatigue, or balance have not been fully examined and will be important to consider when planning any controlled trials of medical cannabis.

Prior studies have shown that age of onset of PD and high income and higher education levels are correlated with CAM use. Our study did not confirm these correlations, perhaps due to the very high rate of CAM use amongst the entire population. Our study also found that 88% of participants reported their physician was aware of their CAM therapy. This is in contrast to previous reports of less than half of PD patients reporting CAM use to their physician. It is difficult to draw conclusions on the reason for this difference but this may reflect cultural shifts in the openness and awareness of physicians regarding CAM as well as shifts in patients' perceptions.

Our study has several notable limitations. Refusal rate was not tracked and therefore our estimate of CAM prevalence in the general PD population may be biased. Presumably, patients that do not use CAM would be more likely to refuse the survey and could lead to an overestimation of CAM usage. In addition, despite our efforts to reach out to the community, the demographics of our study population were typical of a tertiary medical center population, with predominantly white men and highly educated patients, and therefore likely do not fully represent the PD population as a whole. Furthermore, patients living in Colorado may be more likely to utilize CAM than in other regions. All data was self-reported and the survey was self-administered in an anonymous fashion, allowing for misinterpretation of survey questions and making confirmation of responses impossible. There is also a risk that some patients who completed the survey through PD support groups may not have had PD although we assume that this would be a rare occurrence. While we asked patients only to indicate their use of CAM if specifically used for PD, responders may have indicated use even if not specifically for PD. It is also likely that patients used CAM for reasons that we did not ask about such as pain. Furthermore, self-reported efficacy is prone to overreporting of improvement and placebo effect and cannot substitute for data from randomized clinical trials.

Considering the continued growth of CAM use in the general population and amongst PD populations, further randomized clinical trials regarding efficacy of CAM would help physicians guide patients in best use of CAM therapy. Future studies should ideally be placebo-controlled given findings of up to 50% positive placebo effects in prior trials in PD patients [34].

## Supplementary Material

Go to:

Supplementary Material: An electronic copy of the complete survey used in our study is provided here as it was presented to patients.

[Click here to view.](#) (3.8M, pdf)

## Acknowledgment

Go to:

The authors are grateful to Lisa Corbin, MD, for her help with editing the survey and preparing presentations to groups.

## Conflict of Interests

Go to:

The authors declare that there is no conflict of interests regarding the publication of this paper.

## Authors' Contribution

Go to:

T. A. Finseth, J. L. Hedeman, and B. M. Kluger contributed to drafting and revision of the paper. Authors involved in patient recruitment and data collection included K. I. Johnson, M. S. Binder, and B. M. Kluger. Study design was developed by K. I. Johnson and B. M. Kluger. Statistical analysis was performed by R. P. Brown II and B. M. Kluger.

## References

Go to:

1. NCCAM. Complementary, Alternative, or Integrative Health: What's in a name? <http://nccam.nih.gov/health/whatiscam>.
2. Harris P. E., Cooper K. L., Relton C., Thomas K. J. Prevalence of complementary and alternative medicine (CAM) use by the general population: a systematic review and update. *International Journal of Clinical Practice*. 2012;66(10):924–939. doi: 10.1111/j.1742-1241.2012.02945.x. [\[PubMed\]](#) [\[Cross Ref\]](#)
3. Herman P. M., Craig B. M., Caspi O. Is complementary and alternative medicine (CAM) cost-effective? A systematic review. *BMC Complementary and Alternative Medicine*. 2005;5, article 11 [\[PMC free article\]](#) [\[PubMed\]](#)
4. Wang Y., Xie C.-L., Wang W.-W., et al. Epidemiology of complementary and alternative medicine use in patients with Parkinson's disease. *Journal of Clinical Neuroscience*. 2013;20(8):1062–1067. doi: 10.1016/j.jocn.2012.10.022. [\[PubMed\]](#) [\[Cross Ref\]](#)
5. Rajendran P. R., Thompson R. E., Reich S. G. The use of alternative therapies by patients with Parkinson's disease. *Neurology*. 2001;57(5):790–794. doi: 10.1212/WNL.57.5.790. [\[PubMed\]](#) [\[Cross Ref\]](#)
6. Kluger B., Triolo P., Jones W., Jankovic J. The therapeutic potential of cannabinoids for movement disorders. *Movement Disorders*. 2015 doi: 10.1002/mds.26142. [\[PMC free article\]](#) [\[PubMed\]](#) [\[Cross Ref\]](#)
7. Venderová K., Ružička E., Vorříššek V., Višňovský P. Survey on cannabis use in Parkinson's disease: subjective improvement of motor symptoms. *Movement Disorders*. 2004;19(9):1102–1106. doi: 10.1002/mds.20111. [\[PubMed\]](#) [\[Cross Ref\]](#)

8. Reinerman C., Nunberg H., Lanthier F., Heddleston T. Who are medical marijuana patients? Population characteristics from nine California assessment clinics. *Journal of Psychoactive Drugs*. 2011;43(2):128–135. doi: 10.1080/02791072.2011.587700. [\[PubMed\]](#) [\[Cross Ref\]](#)
9. Olanow C. W., Watts R. L., Koller W. C. An algorithm (decision tree) for the management of Parkinson's disease (2001): treatment guidelines. *Neurology*. 2001;56(11) supplement 5:S1–S88. [\[PubMed\]](#)
10. Harris P. A., Taylor R., Thielke R., Payne J., Gonzalez N., Conde J. G. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *Journal of Biomedical Informatics*. 2009;42(2):377–381. doi: 10.1016/j.jbi.2008.08.010. [\[PMC free article\]](#) [\[PubMed\]](#) [\[Cross Ref\]](#)
11. Galicia-Connolly E., Adams D., Bateman J., et al. CAM use in pediatric neurology: an exploration of concurrent use with conventional medicine. *PLoS ONE*. 2014;9(4) doi: 10.1371/journal.pone.0094078.e94078. [\[PMC free article\]](#) [\[PubMed\]](#) [\[Cross Ref\]](#)
12. Stoll S. S., Nieves C., Tabby D. S., Schwartzman R. Use of therapies other than disease-modifying agents, including complementary and alternative medicine, by patients with multiple sclerosis: a survey study. *The Journal of the American Osteopathic Association*. 2012;112(1):22–28. [\[PubMed\]](#)
13. McConnell B. V., Applegate M., Keniston A., Kluger B., Maa E. H. Use of complementary and alternative medicine in an urban county hospital epilepsy clinic. *Epilepsy & Behavior*. 2014;34:73–76. doi: 10.1016/j.yebeh.2014.03.011. [\[PubMed\]](#) [\[Cross Ref\]](#)
14. Wells R. E., Bertisch S. M., Buettner C., Phillips R. S., McCarthy E. P. Complementary and alternative medicine use among adults with migraines/severe headaches. *Headache*. 2011;51(7):1087–1097. doi: 10.1111/j.1526-4610.2011.01917.x. [\[PMC free article\]](#) [\[PubMed\]](#) [\[Cross Ref\]](#)
15. Erwin Wells R., Phillips R. S., McCarthy E. P. Patterns of mind-body therapies in adults with common neurological conditions. *Neuroepidemiology*. 2011;36(1):46–51. doi: 10.1159/000322949. [\[PMC free article\]](#) [\[PubMed\]](#) [\[Cross Ref\]](#)
16. Barnes P. M., Bloom B., Nahin R. L. Complementary and alternative medicine use among adults and children: United States, 2007. *National Health Statistics Reports*. 2009;(12):1–23. [\[PubMed\]](#)
17. Peregoy J. A., Clarke T. C., Jones L. I., Stussman B. J., Nahin R. L. Regional variation in use of complementary health approaches by U.S. adults. *NCHS Data Brief*. 2014;(146):1–8. [\[PubMed\]](#)
18. Kim S. R., Lee T. Y., Kim M. S., Lee M. C., Chung S. J. Use of complementary and alternative medicine by Korean patients with Parkinson's disease. *Clinical Neurology and Neurosurgery*. 2009;111(2):156–160. doi: 10.1016/j.clineuro.2008.09.011. [\[PubMed\]](#) [\[Cross Ref\]](#)
19. Pecci C., Rivas M. J., Moretti C. M., et al. Use of complementary and alternative therapies in outpatients with Parkinson's disease in Argentina. *Movement Disorders*. 2010;25(13):2094–2098. doi: 10.1002/mds.23235. [\[PubMed\]](#) [\[Cross Ref\]](#)
20. Tan L. C. S., Lau P.-N., Jamora R. D. G., Chan E. S. Y. Use of complementary therapies in patients with Parkinson's disease in Singapore. *Movement Disorders*. 2006;21(1):86–89. doi: 10.1002/mds.20662. [\[PubMed\]](#) [\[Cross Ref\]](#)
21. Parkinson Study Group. Effects of tocopherol and deprenyl on the progression of disability in early Parkinson's disease. *The New England Journal of Medicine*. 1993;328(3):176–183. doi: 10.1056/nejm199301213280305. [\[PubMed\]](#) [\[Cross Ref\]](#)
22. Beal M. F. A randomized clinical trial of high-dosage coenzyme q10 in early parkinson disease: no evidence of

benefit. *JAMA Neurology*. 2014;71(5):543–552. doi: 10.1001/jamaneurol.2014.131. [PubMed] [Cross Ref]

23. Degenhardt L., Hall W., Lynskey M. Exploring the association between cannabis use and depression. *Addiction*. 2003;98(11):1493–1504. doi: 10.1046/j.1360-0443.2003.00437.x. [PubMed] [Cross Ref]

24. Ghaffari B. D., Kluger B. Mechanisms for alternative treatments in Parkinson's disease: acupuncture, tai chi, and other treatments. *Current Neurology and Neuroscience Reports*. 2014;14(6) doi: 10.1007/s11910-014-0451-y. [PubMed] [Cross Ref]

25. Li F., Harmer P., Liu Y., et al. A randomized controlled trial of patient-reported outcomes with tai chi exercise in Parkinson's disease. *Movement Disorders*. 2014;29(4):539–545. doi: 10.1002/mds.25787. [PMC free article] [PubMed] [Cross Ref]

26. Paterson C., Allen J. A., Browning M., Barlow G., Ewings P. A pilot study of therapeutic massage for people with Parkinson's disease: the added value of user involvement. *Complementary Therapies in Clinical Practice*. 2005;11(3):161–171. doi: 10.1016/j.ctcp.2004.12.008. [PubMed] [Cross Ref]

27. Stevenson C., Ernst E. Risks associated with spinal manipulation. *The American Journal of Medicine*. 2002;112(7):566–571. doi: 10.1016/s0002-9343(02)01068-9. [PubMed] [Cross Ref]

28. Venderová K., Ružička E., Voršíšek V., Višňovský P. Survey on cannabis use in Parkinson's disease: subjective improvement of motor symptoms. *Movement Disorders*. 2004;19(9):1102–1106. doi: 10.1002/mds.20111. [PubMed] [Cross Ref]

29. Carroll C. B., Bain P. O., Teare L., et al. Cannabis for dyskinesia in Parkinson disease: a randomized double-blind crossover study. *Neurology*. 2004;63(7):1245–1250. doi: 10.1212/01.wnl.0000140288.48796.8e. [PubMed] [Cross Ref]

30. Lotan I., Treves T. A., Roditi Y., Djaldetti R. Cannabis (Medical Marijuana) treatment for motor and non-motor symptoms of parkinson disease: an open-label observational study. *Clinical Neuropharmacology*. 2014;37(2):41–44. doi: 10.1097/wnf.0000000000000016. [PubMed] [Cross Ref]

31. Frankel J. P., Hughes A., Lees A. J., Stern G. M. Marijuana for Parkinsonian tremor. *Journal of Neurology Neurosurgery and Psychiatry*. 1990;53(5, article 436) [PMC free article] [PubMed]

32. Koppel B. S., Brust J. C. M., Fife T., et al. Systematic review: efficacy and safety of medical marijuana in selected neurologic disorders—report of the guideline development subcommittee of the American Academy of Neurology. *Neurology*. 2014;82(17):1556–1563. doi: 10.1212/wnl.0000000000000363. [PMC free article] [PubMed] [Cross Ref]

33. Zuardi A. W., Crippa J. A. S., Hallak J. E. C., et al. Cannabidiol for the treatment of psychosis in Parkinsons disease. *Journal of Psychopharmacology*. 2009;23(8):979–983. doi: 10.1177/0269881108096519. [PubMed] [Cross Ref]

34. Goetz C. G., Wuu J., McDermott M. P., et al. Placebo response in Parkinson's disease: comparisons among 11 trials covering medical and surgical interventions. *Movement Disorders*. 2008;23(5):690–699. doi: 10.1002/mds.21894. [PubMed] [Cross Ref]