

Clinical Outcome Data of First Cohort of Chronic Pain Patients Treated With Cannabis-Based Sublingual Oils in the United Kingdom: Analysis From the UK Medical Cannabis Registry

The Journal of Clinical Pharmacology 2021, 61(12) 1545–1554 © 2021 The Authors. The Journal of Clinical Pharmacology published by Wiley Periodicals LLC on behalf of American College of Clinical Pharmacology DOI: 10.1002/jcph.1961

Michal Kawka, BSc (Hons)¹, Simon Erridge, MBBS, BSc (Hons)^{1,2}, Carl Holvey, MPharm, MRPharmS, DipCLinPharm, IPresc, MBA², Ross Coomber, BSc (Hons), MBBS, FRCS, PGCERT^{2,3}, Azfer Usmani, MBBS, FRCA, FCARCSI, FFPMRCA^{2,4}, Mohammad Sajad, MA, MBBS, FRCA, FFPMRCA^{2,5}, Michael W. Platt, MA, MBBS, FRCA, FFPMRCA^{1,2}, James J. Rucker, MBBS, BSc, MRCPsych, PhD^{2,6,7}, and Mikael H. Sodergren, MBChB (Hons), DIC, PhD, FRCS^{1,2}

Abstract

Cannabis-based medicinal products (CBMPs) are an emerging therapeutic option in the management of primary chronic pain, using the role of the endocannabinoid system in modulating central and peripheral pain processes. Despite promising preclinical data, there is a paucity of high-quality evidence to support the use of CBMPs for chronic pain. This study aimed to investigate the health-related quality-of-life outcomes of patients with chronic pain who were prescribed CBMP oil preparations (Adven, Curaleaf International, Guernsey, UK).

This study is a case series of patients from the UK Medical Cannabis Registry, who were treated with CBMP oils for an indication of chronic pain. The primary outcomes were the changes in Brief Pain Inventory short form, Short-Form McGill Pain Questionnaire-2, Visual Analog Scale Pain, General Anxiety Disorder-7, Sleep Quality Scale, and EQ-5D-5L, at 1, 3, and 6 months. One hundred ten patients were included. Significant improvements in Sleep Quality Scale, EQ-5D-5L pain and discomfort subscale, and Brief Pain Inventory Interference Subscale (P < .05) at 1, 3, and 6 months were demonstrated. There were no notable differences between cannabis-naïve and previous cannabis users in quality-of-life outcomes. The adverse event incidence was 30.0%, with most (n = 58; 92.1%) adverse events being mild or moderate in intensity. Treatment of chronic pain with Adven CBMP oils was associated with an improvement in pain-specific outcomes, health-related quality of life, and self-reported sleep quality. Relative safety was demonstrated over medium-term prescribed use. While these findings must be treated with caution considering the limitations of study design, they can inform future clinical trials.

Keywords

cannabinoids, chronic pain, health-related quality-of-life, medical cannabis

Chronic pain is defined as persistent or recurring pain lasting longer than 3 months, characterized by physical pain, disability, emotional disturbance, and social withdrawal.¹ The global burden of chronic pain is sizable, with an estimated global incidence of 20%.^{2,3} In the United Kingdom alone, 28 million people are thought to be affected by chronic pain.² Chronic pain can be accompanied by depression, anxiety, and sleeping difficulties, which contribute to decreased quality of life and increased incidence of suicide.^{4,5} In addition to the costs to the individual, chronic pain is associated with reduced work productivity, absenteeism, and unemployment. The cost to the UK economy from chronic back pain was estimated at £12.3 billion in 2000 (£21.2 billion, at 2021 inflation levels), with associated costs projected to have risen secondary to an aging population.⁶

While there is increasing recognition of the role of nonpharmacological therapies in the management of ¹Imperial College Medical Cannabis Research Group, Imperial College London, London, UK

²Sapphire Medical Clinics, London, UK

⁴Dartford and Gravesham NHS Trust, Kent, UK

⁵Dudley Group of Hospitals NHS Trust, Dudley, UK

⁶Department of Psychological Medicine, Institute of Psychiatry Psychol-

ogy & Neuroscience, Kings College London, London, UK

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Submitted for publication 14 July 2021; accepted 27 August 2021.

Corresponding Author:

Mikael H. Sodergren, MBChB, DIC, PhD, FRCS, Department of Surgery & Cancer, Imperial College London, Academic Surgical Unit, 10th Floor QEQM, St Mary's Hospital, South Wharf Road, London W2 INY, UK Email: m.sodergren@imperial.ac.uk

³St George's Hospital NHS Trust, London, UK

⁷South London & Maudsley NHS Foundation Trust, London, UK

chronic pain, pharmacological therapies continue to be the mainstay of treatment, despite a paucity of highquality evidence to support their use.^{7–9} For example, while opioid analgesics increased in usage in the early part of the 21st century, there is limited evidence to support their use in chronic pain.^{10–12} This is despite increasing recognition of both short-term adverse effects and the associated risks of long-term use, including addiction, physical dependence, and increased risk of psychological comorbidity.^{11,13,14} Nonsteroidal antiinflammatory drugs, which are commonly prescribed in the setting of musculoskeletal pain, similarly have an unclear role in chronic pain considering their association with gastrointestinal, cardiovascular, and renal adverse events.^{15,16} Gabapentinoids, originally developed for the treatment of epilepsy, have been used with increased frequency off-label for the treatment of chronic pain, partially driven by a desire to find a safer alternative to opioids.^{17,18} However, similar to opioids and nonsteroidal anti-inflammatory drugs, there is a paucity of evidence to support its current breadth of use in clinical practice, in addition to growing evidence of associated harms with chronic use, subversion, and abuse.18-20

In view of the challenges in managing chronic pain, there is growing interest in the endocannabinoid system as a drug target due to its role in modulating central and peripheral pain processes. As such, cannabis-based medicinal products (CBMPs) are an emerging therapeutic option in the management of primary chronic pain of undefined origin, as well as chronic cancer pain and neuropathic pain.^{21–23} CBMPs are a heterogeneous group of pharmaceuticals available as either isolate formulations of cannabinoids, or as broad-spectrum compounds containing other compounds from the cannabis flower with potential therapeutic properties, including terpenes, terpenoids, and flavonoids.²⁴ The 2 cannabinoids to which prescriptions are titrated and with the greatest preliminary research are Δ^9 -tetrahydrocannabinol (THC) and cannabidiol (CBD).²⁴ THC is a partial agonist of the primary receptors of the endocannabinoid system, cannabinoid receptors type 1 and 2 (CB1 and CB2 receptors).^{25,26} CB1 receptors, located in the central nervous system, modulate glutamate and γ aminobutyric acid neurotransmission.^{27,28} CB1 agonism by THC is thought to induce centrally acting analgesic effects, in addition to the psychotropic effects commonly associated with cannabis consumption.^{27,28} CB2 receptors, which are expressed in peripheral immune cells, modulate inflammatory cytokines.^{27,28} However, the extent to which this plays a role in nociception and pain processing is not clear. CBD has opposing effects at cannabinoid receptors. Its primary action is via inhibition of the enzyme fatty acid amino hydrolase, which typically breaks down endogenous cannabinoid receptor agonists (anandamide and 2arachidonoylglycerol).^{29–31} Consequently, CBD leads to increased activation of CB1 and CB2. In clinical studies, CBD and fatty acid amino hydrolase inhibitors appear to have similar clinical effects.^{32,33} However, CBD is also a negative allosteric modulator of CB1 receptors reducing the overall effects of THC and other agonists.³⁴ THC, CBD, and other cannabinoids also act at other receptors implicated in pain pathways, including opioid, transient receptor potential cation channel subfamily V member 1, and serotonin 5-HT3 receptors, each potentially contributing to the overall clinical effects.³⁵

Despite promising preclinical data, there is a paucity of high-quality evidence to support the use of CBMPs. The evidence base, while broad, is inconclusive, variable across chronic pain types, and thus insufficient to inform guidelines, funders, and licensing agencies. For cancer-related pain, placebo-controlled trials have shown clinically significant reduction in pain scores in those using nabiximols, an oromucosal spray containing THC and CBD in a 1.1:1.0 ratio. However, a large trial (n = 397) has shown the difference in effect to be nonsignificant.^{36–38} Systematic reviews of nabiximols in chronic non-cancer and neuropathic pain and CBMPs in musculoskeletal pain have also been inconclusive.^{39,40} However, clinical trials in CBMPs to date have largely been underpowered, performed in acute settings, and failed to account for the heterogeneity of available CBMPs.30,41 Observational studies could be complementary to clinical trials in this field, as they provide insights across a broad spectrum of medicines to guide clinical trials and practice while trial results are awaited.

The UK Medical Cannabis Registry, set up in 2019, captures longitudinal data of patients prescribed CBMPs in the United Kingdom outside of the National Health Service (NHS) and has published outcomes related to health-related quality of life (HRQoL) across all conditions.⁴² Following legalization of medical cannabis in the United Kingdom in 2019, many doctors have preferred sublingual oil as a mode of administration of CBMPs. This study aims to investigate the safety and clinical outcomes of patients with chronic pain enrolled in the UK Medical Cannabis Registry who were treated with cannabis-based medium-chain triglyceride oils (Adven, Curaleaf International, Guernsey, UK).

Methods

Study Overview

This study is an uncontrolled case series of patients identified from the UK Medical Cannabis Registry, and

reported in line with the Strengthening the Reporting of Observational Studies in Epidemiology statement.⁴³ In accordance with the NHS Health Research Authority and Research Ethics Committees guidance, this study did not require formal ethical approval. All participants completed formal, written consent before enrollment in the registry.

Setting and Participants

The UK Medical Cannabis Registry was established in December 2019 and is the first prospective registry launched in the United Kingdom, capturing pseudonymized data on patients treated with CBMPs. It is privately owned and managed by Sapphire Medical Clinics. To date, it is the only clinic that enrolls patients into the registry. Patients are recruited from the United Kingdom and the Channel Islands. Clinicopathological information, comorbidities, drug and alcohol history, and medication information were collected prospectively by clinical staff. Patient-reported outcome measures (PROMs) and adverse event questionnaires were electronically administered to patients at baseline, 1 month, 3 months, and 6 months and the 6 monthly intervals thereafter.

Patient and Data Selection

For this analysis, data were extracted for the initial participants of the UK Medical Cannabis Registry who were prescribed cannabis-based oil preparations (Adven) for the indication of chronic pain and had recorded PROMs at baseline with at least 1 follow-up datum (1, 3, and/or 6 months). In the United Kingdom, prescriptions of CBMPs are only prescribed once other treatments have proven ineffective or inappropriate. Only patients who were prescribed exclusively Adven CBMPs in the form of oil preparation and no other CBMPs and modes of administration were included in this analysis. Data regarding demographic details, including age, sex, and occupation, were recorded. Participant body mass index (BMI) was also extracted. Data on the relevant comorbidities contributing to the Charlson Comorbidity Index, a widely used prognostic scoring model for 10-year mortality, was collected and a score calculated for each patient.44 While not wholly representative of typical comorbidities that accompany chronic pain, the Charlson Comorbidity Index was chosen to be collected across the UK Medical Cannabis Registry, as it is the most commonly used comorbidity scoring system used in registry studies, allowing for direct comparison of comorbid status between cohorts.45

Drug and alcohol data on patients were extracted and analyzed, including smoking status, smoking packyears, alcohol units per week, and cannabis status. For those who had previously or were presently taking nonprescription cannabis, a novel metric of "gram years" was calculated as previously described by our group.⁴² All CBMP prescriptions were recorded and analyzed, including company, formulation, method of administration, CBD concentration, THC concentration, and strain.

All participants are administered quality-of-life PROMs questionnaires, including EQ-5D-5L, General Anxiety Disorder-7 (GAD-7), and Single-Item Sleep Quality Scale (SQS). Fibromyalgia patients were administered the Fibromyalgia Severity Scale, while all patients with other chronic pain etiologies were administered the Visual Analog Scale (VAS) Pain, Brief Pain Inventory Short Form (BPI), and Short-Form McGill Pain Questionnaire-2.⁴⁶⁻⁵¹

The BPI is a validated scale that assesses pain at its "worst," "least," "average," and "now" (current pain) to produce a severity score from 0 to 10, as well as measuring interference score, which measures how much pain has interfered with 7 daily activities, including general activity, walking, work, mood, enjoyment of life, relations with others, and sleep.

The McGill Pain Questionnaire includes painspecific descriptors, which patients score the intensity of, as a number from 1 to 5, in which each number is associated with the following words: 1, "mild"; 2, "discomforting"; 3, "distressing"; 4, "horrible"; and 5, "excruciating." The descriptors fall into 5 major groups: continuous, intermittent, affective, neuropathic, and overall, for which the total scores are calculated.

The Fibromyalgia Severity Scale is derived from the 2016 diagnostic criteria for fibromyalgia.⁵¹ It ranges from 0 to 31 and is a sum of the widespread pain index, a value measuring the areas of pain experienced by a patient with fibromyalgia, and the symptom severity score, a measure of the severity of symptoms associated with fibromyalgia including fatigue, waking feeling unrefreshed, cognitive symptoms, headaches, lower abdominal pain, and depression.

The EQ-5D-5L is a 2-part tool that measures the quality of life across 5 domains (mobility, self-care, usual activities, pain or discomfort, anxiety or depression) with 5 levels of severity (no problems, slight problems, moderate problems, severe problems, and extreme problems). These 5 domains and levels correspond to 1 of 3125 health states, which are mapped to EQ-5D-5L index values according to a technique described by van Hout et al.⁵² An index value of <0 represents a health state worse than death, while a score of 1 indicates perfect HRQoL. This is the preferred method of measuring HRQoL by the National Institute for

Health and Care Excellence.⁵³ The second part of the EQ-5D-5L is the EQ-VAS, which consists of a vertical scale of 0 to 100, whereby "100" corresponds to the "best health you can imagine" and "0" corresponds to the "worst health you can imagine."⁴⁶

For the GAD-7 score, registry participants are asked about how often over the past 2 weeks they had been bothered by the core symptoms of generalized anxiety disorder, generating a score from 0 to 21, where a higher score is consistent with worse anxiety symptoms.⁴⁷

The SQS is a validated question of sleep quality over the past 7 days only with sleep quality rated from 0 to 10, wherein "10" signifies "excellent" and "0" denotes "terrible."⁴⁸

Patients were also asked to rate their pain on a VAS of 0 to 10, where "0" is "no pain at all" and "10" is the "worst pain that they can imagine."

Participants reported adverse events at 1, 3, and 6 months from baseline, either through self-reporting or during routine follow-up with a clinician. Adverse events were recorded in accordance with the common terminology criteria for adverse events version 3.0.⁵⁴

Outcomes

Primary outcome measures for this analysis were changes in BPI, Short-Form McGill Pain Questionnaire-2, VAS Pain, GAD-7, SQS, and EQ-5D-5L PROMs, at 1, 3, and 6 months. Secondary outcomes included analysis of adverse effects in terms of severity and incidence.

Statistical Analysis

Data from PROMs were analyzed compared to baseline at 1, 3, and 6 months. Normality was tested via a Shapiro-Wilk test. Parametric data were presented as a mean \pm standard deviation, while nonparametric data were presented as median (interquartile range [IQR]). A preplanned subgroup analysis comparing patients who have self-identified as active cannabis users before starting CBMP therapy with a subgroup composed of cannabis-naïve and ex-cannabis users was conducted. Baseline PROM data and change scores (calculated as a difference between baseline and follow-up data) were compared between subgroups. An adverse events profile was also analyzed.

Statistical analysis was performed with a *t*-test or Wilcoxon rank-sum test depending on whether the data were parametric or nonparametric, respectively. Statistical significance was defined using *P* value <.05. (R version 4.0.3; R Foundation for Statistical Computing, Vienna, Austria) was used for data visualization and analysis.

Table 1. Participants' Demographic and Clinical Characteristics

Baseline Characteristics	No. (%)/Mean \pm SD
Age, y	52.I ± 15.4
Female	56 (50.90)
Male	54 (49.10)
Body mass index, kg/m ²	$\textbf{28.00} \pm \textbf{5.56}$
Occupation	
Elementary occupations	5 (4.55)
Professional	19 (17.27)
Retired	7 (6.36)
Unemployed	34 (30.91)
Other occupation	41 (37.27)
Undisclosed	4 (3.64)
Cannabis status	
Never used	71 (64.55)
Current user	26 (23.64)
Ex-user	13 (11.81)
Smoking status	
Never smoked	55 (50.00)
Current smoker	14 (12.72)
Ex-smoker	41 (37.27)
Charlson Comorbidity Index	1.69 ± 1.94
Mean alcohol consumption per week, units	$\textbf{6.39} \pm \textbf{10.51}$
Mean pack-years	11.96 \pm 11.72
Mean cannabis lifetime use, gram years	$\textbf{9.44} \pm \textbf{24.58}$

SD, standard deviation.

Results

Data extraction included the first 831 patients who had been registered on the UK Medical Cannabis Registry. When restricted to patients who had received treatment for >1 month, 737 patients remained. Out of 737 patients, 449 had received CBMPs for chronic pain (primary diagnosis) and 257 received only Adven oil preparation, with no other CBMPs prescribed. From these, 148 had completed baseline PROM data and 110 of those had completed at least 1 follow-up data point. Of these, 100 patients had recorded PROMs at 1 month, while 54 patients had recorded PROMs at 3 months and 20 patients had PROMs at 6 months.

Demographic details are presented in Table 1. The mean age of patients was 52.07 (\pm 15.43). Nearly half (n = 54; 49.1%) of the patients were female. The mean BMI of participants was 28.02 (\pm 5.66). The majority of the patients have never used cannabis (n = 71; 64.55%), with nearly a quarter being current users (n = 26; 23.64%) and the rest being ex-users (n = 13; 11.81%). The mean cannabis lifetime use was 9.44 gram-years.

Table 2 outlines the primary diagnosis for which treatment was initiated. The most common primary diagnosis was chronic noncancer pain (n = 53; 48.2%), followed by neuropathic pain (n = 26; 23.6%) and fibromyalgia (n = 18; 16.3%). Fifty-two (47.3%) and 14 (12.7%) patients, respectively, also had a secondary or tertiary indication for CBMP therapy.

Table 2. Distribution of Indications for CBMP Therapy

Diagnosis	Primary, n (%)	Secondary, n (%)	Tertiary, n (%)
Chronic noncancer pain	53 (48.2)	18 (13.4)	2 (1.8)
Neuropathic pain	26 (23.6)	7 (6.3)	2 (1.8)
Fibromyalgia	18 (16.2)	12 (10.9)	0 (0.0)
Ehlers-Danlos	6 (5.4)	5 (4.5)	I (0.9)
Complex regional pain syndrome	4 (5.4)	0 (0.0)	0 (0.0)
Cancer pain	3 (2.7)	0 (0.0)	0 (0.0)
Insomnia	0 (0.0)	3 (2.7)	0 (0.0)
Migraine	0 (0.0)	2 (1.8)	3 (2.7)
Depression	0 (0.0)	I (0.9)	2 (1.8)
Agoraphobia	0 (0.0)	0 (0.0)	I (0.9)
Anxiety	0 (0.0)	2 (1.8)	0 (0.0)
Autism spectrum disorder	0 (0.0)	2 (1.8)	I (0.9)
Eating disorder	0 (0.0)	0 (0.0)	I (0.9)
PTSD	0 (0.0)	0 (0.0)	l (0.9)

CBMP, cannabis-based medicinal product; PTSD, posttraumatic stress disorder.

CBMP Dosing and Mode of Administration

The majority of patients had 2 cannabis-based oil preparations prescribed (n = 105; 95.5%), with the rest of patients being prescribed a single CBMP (n = 5; 4.5%). Two hundred five CBMPs were prescribed across the whole cohort. Sativa strains were the most commonly prescribed (n = 109; 53.2%), followed by hybrid strains (n = 98; 47.8%). Fifty-nine patients (53.6%) were prescribed the same combination of 20 mg of CBD oil (sativa strain) and 1 mg of THC oil (hybrid strain). The median CBD dose at baseline was 20.0 mg per day (IQR, 20.0-20.0 mg), while the median THC dose at baseline was 1.00 mg per day (IQR 1.0-2.0 mg). The majority of oils were prescribed for oral admission (n = 167; 81.5%).

Patient-Reported Outcome Measures

Table 3 outlines full paired results from baseline to 1 month, 3 months, and 6 months (where data are available). Statistically significant improvements in HRQoL were demonstrated at 1, 3, and 6 months in SQS, EQ-5D-5L pain and discomfort subscale, and Brief Pain Inventory Interference Subscale (P < .05). Statistically significant improvements were also demonstrated at 1 and 3 months only in the EQ-5D-5L Index, VAS Pain, and EQ-5D-5L VAS (P < .05). It is worth noting that due to the bimodal distribution of the data, some comparisons were significant, despite the median baseline score and follow-up score being the same.

Effect of Previous Cannabis Use

The majority of current users declared using cannabis daily (n = 24; 88.5%), with median daily usage of 0.75 g (IQR, 0.35-2.00 g). On subgroup analysis, no statistical significance difference between cannabis-naïve and current users was found at baseline for all the PROMs (P > .05). When the change scores were considered,

1524604, 2021, 12, Downloaded from https://accp1.ninelibrary.wiey.com/doi/10.102/jcph.1961, Wiely Online Library on [2406/2025]. See the Terms and Conditions (https://anlinelibrary.wiely.com/terms-and-conditions) on Wiely Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

the cannabis-naïve group had a significantly larger difference in scores at 1 month (median, -0.52 vs + 0.22; P = .044). Moreover, at 3-month follow-up, the current cannabis user subgroup had a larger difference in McGill's Continuous Pain Subscale scores (median, -3.42 vs 0.50, P = .036). Full subgroup analysis results can be found in Table S1.

Adverse Events

Reported adverse events are described in full in Table 4. There were 63 total reported adverse events reported by 33 patients (30.0%). The most commonly experienced adverse events were nausea (n = 9; 14.3%) and dizziness (n = 7; 11.1%). The majority of adverse events were mild or moderate (n = 58; 92.1%), with only 2 events being described as severe and 1 as disabling. Only 1 of the 63 adverse events (1.6%) was reported by a current cannabis user, with the rest being reported by patients who were either ex-users (n = 4, 6.3%) or cannabis naïve (n = 58; 92.1%).

Discussion

This analysis presents the short-term outcomes of the first cohort of patients prescribed CBMP oils for chronic pain in the United Kingdom. Significant improvements in HRQoL were demonstrated in SQS, EQ-5D-5L pain and discomfort subscale, and Brief Pain Inventory Interference Subscale (P < .05) at all followup points recorded to date (at 1, 3, and 6 months). There were no notable differences between cannabis-naïve and previous cannabis users in terms of quality-oflife outcomes. The adverse event incidence was 30.0%, with the majority of adverse events being either mild or moderate in intensity (n = 58; 92.1%).

Registries have already been established for monitoring of CBMP prescriptions in chronic pain globally. A recent analysis of German pain e-Registry in 2017, which investigated the effectiveness and safety of THC/CBD spray for chronic pain management in a cohort of 800 patients found that 56% of patients had >50% improvement in >5 of 9 aggravated symptom relief-9 scale domains. These included improvements in pain intensity (67.5%), depression (66.5%), overall well-being (61.3%), anxiety (57.6%), disabilities in daily life (56.3%), sleep (47.0%), and physical (42.1%) and mental quality-of-life (17.4%).55 This is in line with the findings of the present study, in terms of improvements in pain intensity, overall HRQoL, and sleep quality, all of which were significant at 1, 3, and 6 months in our analysis. However, in the present study, there was no detectable improvement in the EQ-5D-5L domains of mobility, self-care, and usual activities. This may suggest that CBMP oils may be associated with improved perception of pain without resulting

Table 3	. Paired	Baseline and	Follow-Up	Patient-Reported	Outcome	Measures
---------	----------	--------------	-----------	------------------	---------	----------

PROM	Follow-Up	n	Scores at Baseline	Scores at Follow-Up	Р
					· · ·
GAD-7	l mo	98	4.00 (0.20-8.00)	2.50 (0.00-8.00)	.346
	3 mo	51	4.00 (0.50-9.00)	3.00 (1.00-8.00)	.663
	6 mo	20	5.00 (1.00-11.80)	6.00 (2.50-12.00)	.914
SQS	l mo	88	5.06 ± 2.49	5.77 ± 2.33	.001**
	3 mo	47	4.83 ± 2.42	6.09 ± 2.47	<.001 ^{%%%}
	6 mo	14	4.57 ± 2.34	6.29 ± 2.30	.024**
Pain VAS	l mo	68	6.37 ± 1.84	5.97 ± 1.71	.016*
	3 mo	41	6.61 ± 1.64	5.56 ± 2.14	<.001 ^{solok}
	6 mo	12	6.33 ± 1.72	5.42 ± 2.50	.190
EQ-5D-5L Mobility	l mo	92	3.00 (2.00-4.00)	3.00 (1.80-3.00)	.200
	3 mo	51	3.00 (2.00-4.00)	3.00 (2.00-4.00)	.655
	6 mo	19	3.00 (2.00-4.00)	3.00 (2.00-3.50)	.964
EQ-5D-5L Self-Care	l mo	92	2.00 (1.00-3.00)	2.00 (1.00-5.00)	.997
	3 mo	51	2.00 (1.00-3.00)	1.00 (1.00-3.00)	.718
	6 mo	19	1.00 (1.00-3.00)	2.00 (1.00-3.00)	.514
EQ-5D-5L Usual Activities	l mo	92	3.00 (2.00-4.00)	3.00 (2.00-3.00)	.073
	3 mo	51	3.00 (2.00-4.00)	2.00 (2.00-3.50)	.120
	6 mo	19	3.00 (2.00-4.00)	3.00 (2.50-3.50)	.575
EQ-5D-5L Pain and Discomfort	l mo	92	3.00 (3.00-4.00)	3.00 (2.00-4.00)	.028*
	3 mo	51	3.00 (3.00-4.00)	3.00 (3.00-4.00)	.009***
	6 mo	19	4.00 (3.00-4.00)	3.00 (2.00-3.50)	.013*
EQ-5D-5L Anxiety and Depression	l mo	92	2.00 (1.00-3.00)	2.00 (1.00-3.00)	.545
	3 mo	51	2.00 (1.00-3.00)	1.00 (1.00-2.00)	.135
	6 mo	19	2.00 (1.00-3.00)	2.00 (1.00-3.50)	.878
EQ-5D-5L Index	l mo	92	$\textbf{0.447} \pm \textbf{0.299}$	0.511 ± 0.290	.005***
	3 mo	51	0.401 ± 0.319	$\textbf{0.496} \pm \textbf{0.299}$.003***
	6 mo	19	$\textbf{0.357} \pm \textbf{0.343}$	$\textbf{0.404} \pm \textbf{0.345}$.295
EQ-5D-5L VAS	l mo	92	$\textbf{52.43} \pm \textbf{21.43}$	$\textbf{56.72} \pm \textbf{21.28}$.028*
	3 mo	50	$\textbf{50.76} \pm \textbf{21.60}$	$\textbf{57.46} \pm \textbf{21.48}$.030*
	6 mo	18	$\textbf{46.17} \pm \textbf{23.55}$	$\textbf{49.83} \pm \textbf{22.88}$.480
Brief Pain Inventory—Severity Subscale	l mo	68	5.09 \pm 1.95	4.75 ± 1.85	.071
	3 mo	36	5.40 ± 1.76	$\textbf{4.37} \pm \textbf{1.91}$.014*
	6 mo	13	$\textbf{5.29} \pm \textbf{2.13}$	4.15 ± 1.40	.012*
Brief Pain Inventory—Interference Subscale	l mo	68	$\textbf{4.95} \pm \textbf{2.12}$	$\textbf{4.38} \pm \textbf{2.26}$.0182*
	3 mo	36	5.24 ± 2.03	$\textbf{4.22} \pm \textbf{2.33}$.029*
	6 mo	13	5.78 ± 2.25	3.96 ± 1.85	<.001***
McGill's Pain Scale—Continuous Pain Subscale	l mo	56	4.42 (2.29-6.4)	3.58 (1.79-5.71)	.255
	3 mo	33	4.33 (2.67-6.67)	3.67 (1.83-5.17)	.213
	6 mo	11	4.50 (2.92-6.08)	2.83 (1.00-5.42)	.211
McGill's Pain Scale—Intermittent Pain Subscale	l mo	56	4.42 (1.33-5.71)	3.42 (1.25-5.04)	.343
	3 mo	33	4.67 (1.67-5.83)	2.83 (0.67-5.17)	.162
	6 mo	11	5.00 (1.17-5.50)	1.50 (0.17-3.83)	.129
McGill's Pain Scale—Affective Pain Subscale	l mo	56	3.62 (1.75-5.75)	2.50 (1.25-4.31)	.124
	3 mo	33	3.75 (2.00-5.00)	2.25 (0.75-4.00)	.057
	6 mo	11	5.00 (2.38-6.00)	1.25 (0.25-4.12)	.030*
McGill's Pain Scale— Overall Score	l mo	56	3.79 (2.22-5.31)	3.09 (1.63-4.71)	.195
	3 mo	33	3.98 (2.36-5.44)	2.94 (1.15-4.67)	.109
	6 mo	11	4.71 (2.58-4.17)	2.04 (0.73-4.77)	.065
McGill's Pain Scale—Neuropathic Pain Subscale	l mo	56	3.08 (1.17-4.37)	2.75 (1.00-3.71)	.316
	3 mo	33	3.33 (1.83-4.67)	3.33 (0.67-4.50)	.445
	6 mo	11	3.67 (3.25-4.17)	2.83 (0.75-3.75)	.293
Fibromyalgia Severity Scale	l mo	12	23.50 (18.50-25.00)	20.50 (15.25-23.75)	.340
	3 mo	9	18.00 (16.00-24.00)	18.00 (13.00-22.00)	.505

GAD-7, General Anxiety Disorder-7; SQS, Sleep Quality Scale; VAS, Visual Analog Scale.

* P < .05, **P < .01, ***P < .001.

Table 4. Adverse Event Profile

	Life Threatening/					
Adverse Events	Severity Mild	Moderate	Severe	Disabling	Unknown	Total, n (%)
Nausea	9	0	0	0	0	9 (14.3)
Dizziness	5	2	0	0	0	7 (11.1)
Headache	2	4	I	0	0	7 (11.1)
Constipation	5	I	0	0	0	6 (9.6)
Fatigue	5	0	0	0	0	5 (7.9)
Dry mouth	4	I	0	0	0	5 (7.9)
Other	I	3	0	0	I	5 (7.9)
Memory impairment	0	4	0	0	0	4 (6.3)
Vomiting	2	0	0	0	0	2 (3.2)
Concentration impairment	I	I	0	0	0	2 (3.2)
Upper abdominal pain	2	0	0	0	0	2 (3.2)
Somnolence (sleepy/ drowsy)	0	2	0	0	0	2 (3.2)
Amnesia	I	0	0	0	0	l (l.6)
Dyspepsia	0	0	I	0	0	l (l.6)
Insomnia	I.	0	0	0	0	l (l.6)
Lethargy	0	I	0	0	0	l (l.6)
Coordination/Balance/Speech impairment	I.	0	0	0	0	l (l.6)
Vertigo (spinning/dizziness)	0	0	0	I	0	l (l.6)
Total, n (%)	39 (61.9)	19 (30.2)	2 (3.2)	l (l.6)	l (l.6)	63 (100)

in functional change. The study was not designed to assess for differences in this subscale, and further evaluation of the role of CBMPs in effecting function in the setting of chronic pain will be required. The improvements in sleep quality, could be secondary to the endocannabinoid system's role in the regulation of the sleep-wake cycle, through modulation of circadian rhythm by acting on the suprachiasmatic nucleus.⁵⁶ Similar improvements in sleep quality in the context of chronic conditions, have further been shown in a recent meta-analysis of 8 placebo-controlled trials.⁵⁷

The present analysis has demonstrated a statistically significant improvement in health-related quality of life in a range of pain-related metrics (EQ-5D-5L pain and discomfort subscale, and Brief Pain Inventory Interference Subscale). This is consistent with the results of a prospective observational study of 751 patients in Canada, who were prescribed medical cannabis for chronic pain conditions, which found a decrease in both pain interference and severity subscale at 12 months.⁵⁸ This is also consistent with the initial analysis of patients from the UK Medical Cannabis Registry, which noted an improvement in HRQoL, as measured by the EQ-5D-5L index value, across all conditions.⁴² A 2017 systematic review and metaanalysis of HRQoL outcomes in those prescribed CBMPs or synthetic cannabinoids for all conditions failed to demonstrate an impact on HRQoL.⁵⁹ While formal analysis was not performed on pain-specific studies, it was notable that pain studies were more likely to have a positive impact on HRQoL. This review was also limited, as HRQoL was a secondary

outcome of most included studies. What is more, the heterogeneity in dose and consistency of cannabis regarding THC:CBD ratio and methods of intake prevents direct comparison between the studies. It should be noted that due to interpersonal differences in response to CBMPs, and variability between cannabis plant strains, a degree of variability between prescribed protocols will inevitably exist and generally limit the generalizability of studies on cannabinoids. In this study, analysis was focused on a single CBMP oil range to limit this variability where possible.

There was no noted improvement in any aspects of McGill's Pain Scale, except the affective pain scale at 6 months. This is in keeping with the latest Cochrane review of cannabis products in adult neuropathic pain, which concluded that there was no clinically significant difference in pain relief in those with neuropathic pain.⁶⁰ This review did find low-quality evidence of a difference in pain intensity and sleep, again corroborated by the present study. However, this study comprises only 35 (31.5%) participants with neuropathic pain, including secondary and tertiary diagnoses, and therefore may not reveal the outcomes for those with pain of a neuropathic origin. Preclinical studies have highlighted a potential role of THC and CBD in neuropathic pain models; however, this is not wholly supported at present by clinical data. In future studies, we aim to assess the effect of CBMPs in select patients from the UK Medical Cannabis Registry with neuropathic pain for better evaluation.⁶¹

While self-reporting of adverse events, which is used in the UK Medical Cannabis Registry can lead to selection bias, the overall adverse event rate in this cohort was 30.0%, which is comparable to the incidence reported in the postmarketing safety analysis of nabiximols by Etges et al $(31.3\%)^{62}$ but higher than in the study by Oreja-Guevara et al (20.0%).⁶³ Interestingly, all but 1 adverse event was reported by patients who have self-reported never previously using cannabis, suggesting potential initial adjustment period in which dosing regimens are being optimized by patients, after which adverse events subside. Moreover, those who use cannabis before starting treatment with CBMPs are subject to selection bias with those with significant side effects unlikely to seek treatment. As the UK cannabis registry is expanded, a more in-depth analyses of adverse events rate over time should be conducted to elucidate the true incidence.

While registry studies provide a resource-efficient method to collect naturalistic data, this study has inherent limitations. Due to a lack of comparator arm or a control group, this introduces biases and limits the ability to make any statements regarding causation for the associations found. What is more, while efforts have been taken to reduce the selection bias, it is not eliminated, as all included patients were prescribed CBMP therapy via a private prescription. This limits access to CBMPs to those who cannot afford the costs associated with such therapy. Selection bias was compounded by the exclusion of patients who did not have baseline data or did not complete follow-up data and must be taken into account in interpreting the results. However, to limit the effect of incomplete or missing data, the study was restricted to those with complete baseline data. While the potential effect of previous cannabis use could have confounded the results, on subgroup analysis we have found no notable differences between cannabis-naïve and previous-user subgroups in terms of PROMs (Table S1). It is also worth noting that regression toward the mean phenomenon could have accounted for some of the differences observed, especially for patients with multiple repeated observations, such as those who had a 1-, 3-, and 6-month follow-up set of PROMs.

While there is heterogeneity in the etiology of chronic pain treated in the context of this study, this is the first study of its kind of patients from the United Kingdom. This study focuses on short-term outcomes, and medium- and long-term efficacy and safety outcomes of CBD/THC oils in the treatment of chronic pain are still unknown, as data were not sufficient to conduct the analysis at 12-month follow-up. Going forward, as more patients are being prescribed CBMPs in the United Kingdom, analyses of long-term outcomes are pertinent. A more in-depth study into specific prescribing protocols (dose, strain, mode of admission) and investigating dose-response relationship are needed for establishing optimal dosing regimens. Interactions and potential synergistic effect of other analgesics with CBMPs should also be explored, as a single agent is unlikely to be used in isolation for chronic pain management.

Conclusion

These results suggest that treatment of chronic pain with selected prescription regimens of sublingual CBMP oils (Adven) is associated with an improvement in pain-specific outcomes in addition to HRQoL and self-reported sleep quality. Moreover, it details the incidence profile of adverse events, particularly highlighting how severe or disabling side effects are rare over medium-term treatment, which is in line with the previously published data. Due to notable limitations of this study, definite conclusions on specific prescription regimen efficacy cannot be drawn; however, the 20-mg CBD: 1-mg THC prescription protocol, which was most commonly used in the study, shows promising initial results. As the UK Medical Cannabis Registry increases in the number of participants and follow-up available, further product-specific analyses are planned across conditions, including oils and flower CBMPs.

Conflicts of Interest

M.K. is a medical student at Imperial College London. He has no shareholdings in pharmaceutical companies. S.E. is a junior doctor and undertakes paid consultancy work at Sapphire Medical Clinics. He is an honorary clinical research fellow at Imperial College London. He has no shareholdings in pharmaceutical companies. C.H. is chief clinical pharmacist at Sapphire Medical Clinics. He has no shareholdings in pharmaceutical companies. R.C. is a consultant orthopedic surgeon, a director, and a shareholder at Sapphire Medical Clinics and a consultant at St George's Hospital, London. He has no shareholdings in pharmaceutical companies. A.U. is a pain specialist at Sapphire Medical Clinics (London) and a consultant at Dartford and Gravesham NHS Trust. He has no shareholdings in pharmaceutical companies. M.S. is a pain specialist at Sapphire Medical Clinics (London) and a consultant at Dudley Group of Hospitals NHS Trust. He has no shareholdings in pharmaceutical companies. M.P. is a consultant in pain services and a director and a shareholder at Sapphire Medical Clinics (London). He has no shareholdings in pharmaceutical companies. J.R. is a consultant psychiatrist, a director, and a shareholder at Sapphire Medical Clinics (London). He is an honorary consultant psychiatrist at the South London & Maudsley NHS Foundation Trust, and a National Institute for Health Research (NIHR) Clinician Scientist Fellow at the Centre for Affective Disorders at King's College London. He is funded by a fellowship (CS-2017-17-007) from the NIHR. He leads the Psychedelic Trials Group at King's College London. King's College London receives grant

funding from COMPASS Pathways PLC to undertake phase 1 and phase 2 trials with psilocybin. COMPASS Pathways PLC has paid for him to attend trial-related meetings and conferences to present the results of research using psilocybin. He has undertaken paid consultancy work for Beckley PsyTech and Clerkenwell Health. Payments for consultancy work are received and managed by King's College London, and he does not benefit personally. He has no shareholdings in pharmaceutical companies. M.S. is a consultant hepatopancreatobiliary surgeon at Imperial College Healthcare NHS Trust, London, and a Senior Clinical Lecturer at Imperial College London. He is a founder, director, and a shareholder at Sapphire Medical Clinics and Research Director at Curaleaf International. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, or the Department of Health.

Data Availability Statement

Data that support the findings of this study are available from the UK Medical Cannabis Registry. Restrictions apply to the availability of these data. Data specifications and applications are available from the corresponding author.

References

- Aziz Q, Barke A, Bennett MI, et al. A classification of chronic pain for ICD-11. *Pain*. 2015;156(6):1003-1007.
- 2. Fayaz A, Croft P, Langford RM, Donaldson LJ, Jones GT. Prevalence of chronic pain in the UK: a systematic review and meta-analysis of population studies. *BMJ Open.* 2016;6(6).
- Gureje O, Von Korff M, Simon GE, Gater R. Persistent pain and well-being. JAMA. 1998;280(2):147.
- Ratcliffe GE, Enns MW, Belik SL, Sareen J. Chronic pain conditions and suicidal ideation and suicide attempts: an epidemiologic perspective. *Clin J Pain*. 2008;24(3):204-210.
- Tang NKY, Crane C. Suicidality in chronic pain: a review of the prevalence, risk factors and psychological links. *Psychol Med.* 2006;36(5):575-586.
- Maniadakis N, Gray A. The economic burden of back pain in the UK. *Pain*. 2000;84(1):95-103.
- Hylands-White N, Duarte R V., Raphael JH. An overview of treatment approaches for chronic pain management. *Rheumatol Int*. 2017;37(1):29-42.
- Ventafridda V, Saita L, Ripamonti C, De Conno F. WHO guidelines for the use of analgesics in cancer pain. *Int J Tissue React*. 1985;7(1):93-96.
- National Institue for Health and Care Excellence. Chronic pain (primary and secondary) in over 16s: assessment of all chronic pain and management of chronic primary pain [NG193]. 2019.
- Jani M, Yimer BB, Sheppard T, Lunt M, Dixon WG. Time trends and prescribing patterns of opioid drugs in UK primary care patients with non-cancer pain: a retrospective cohort study. *PLoS Med.* 2020;17(10):1-16.
- Benyamin R, Trescot AM, Datta S, et al. Opioid complications and side effects. *Pain Physician*. 2008;11(Spec. Iss. 2):105-120.
- Gudin J, Kaufman AG, Datta S. Are opioids needed to treat chronic low back pain? A review of treatment options and analgesics in development. *J Pain Res.* 2020;13:1007-1022.
- Scherrer JF, Salas J, Copeland LA, et al. Prescription opioid duration, dose, and increased risk of depression in 3 large patient populations. *Ann Fam Med.* 2016;14(1):54-62.

1553

5524604, 2021, 12, Downloaded from https://accp1.olinielibrary.wiley com/doi/10.1002/jcph.1961, Wiley Online Library or [2406/2025]. See the Terms and Conditions (https://olinielibrary.wiley com/terms-and-conditions) on Wiley Online Library for rules of use; OA attrices are governed by the applicable Creative Commons License

- Chu LF, Clark DJ, Angst MS. Opioid tolerance and hyperalgesia in chronic pain patients after one month of oral morphine therapy: a preliminary prospective study. *J Pain.* 2006;7(1):43-48.
- Gore M, Tai KS, Sadosky A, Leslie D, Stacey BR. Use and costs of prescription medications and alternative treatments in patients with osteoarthritis and chronic low back pain in community-based settings. *Pain Pract.* 2012;12(7):550-560.
- Machado GC, Abdel-Shaheed C, Underwood M, Day RO. Non-steroidal anti-inflammatory drugs (NSAIDs) for musculoskeletal pain. *BMJ*. 2021;372:1-6.
- 17. Gu X, Chen TC, Su TL, Steinke D, Chen LC. Investigating the prescribing trajectory and geographical drug utilisation patterns of gabapentinoids in primary care in England: an ecological study. *Br J Clin Pharmacol.* 2021;(September 2020):1-12.
- Goodman CW, Brett AS. Gabapentin and pregabalin for pain is increased prescribing a cause for concern? *N Engl J Med.* 2017;377(5):411-13.
- Peckham AM, Ananickal MJ, Sclar DA. Gabapentin use, abuse, and the US opioid epidemic: the case for reclassification as a controlled substance and the need for pharmacovigilance. *Risk Manag Healthc Policy*. 2018;11:109-116.
- Horne AW, Vincent K, Hewitt CA, et al. Gabapentin for chronic pelvic pain in women (GaPP2): a multicentre, randomised, double-blind, placebo-controlled trial. *Lancet*. 2020;396(10255):909-917.
- 21. Lee G, Grovey B, Furnish T, Wallace M. Medical cannabis for neuropathic pain. *Curr Pain Headache Rep.* 2018;22(1).
- Blake A, Wan BA, Malek L, et al. A selective review of medical cannabis in cancer pain management. *Ann Palliat Med.* 2017;6(suppl 2):S215-S222.
- Romero-Sandoval EA, Fincham JE, Kolano AL, Sharpe BN, Alvarado-Vázquez PA. Cannabis for chronic pain: challenges and considerations. *Pharmacotherapy*. 2018;38(6):651-662.
- Hanuš LO, Meyer SM, Muñoz E, Taglialatela-Scafati O, Appendino G. *Phytocannabinoids: A Unified Critical Inventory*. 33; 2016.
- Vučkovic S, Srebro D, Vujovic KS, Vučetic Č, Prostran M. Cannabinoids and pain: New insights from old molecules. *Front Pharmacol.* 2018;9:1-19.
- Paronis CA, Nikas SP, Shukla VG, Makriyannis A. Δ9-Tetrahydrocannabinol acts as a partial agonist/antagonist in mice. *Behav Pharmacol*. 2012;23(8):802-805.
- 27. Ibsen MS, Connor M, Glass M. Cannabinoid CB 1 and CB 2 receptor signaling and bias . *Cannabis Cannabinoid Res.* 2017;2(1):48-60.
- Szabo B, Schlicker E. Effects of cannabinoids on neurotransmission. In: *Handbook of Experimental Pharmacology*. 168; 2005:327-365.
- Turcotte C, Blanchet MR, Laviolette M, Flamand N. The CB2 receptor and its role as a regulator of inflammation. *Cell Mol Life Sci.* 2016;73(23):4449-4470.
- Maharajan MK, Yong YJ, Yip HY, et al. Medical cannabis for chronic pain: can it make a difference in pain management? J Anesth. 2020;34(1):95-103.
- Leweke F, Piomelli D, Pahlisch F, et al. Cannabidiol enhances anandamide signaling and alleviates psychotic symptoms of schizophrenia. *Transl Psychiatry*. 2012;2:e94 https://doi.org/10. 1038/tp.2012.15.
- 32. Bisogno T, Hanuš L, De Petrocellis L, et al. Molecular targets for cannabidiol and its synthetic analogues: Effect on vanilloid VR1 receptors and on the cellular uptake and enzymatic hydrolysis of anandamide. *Br J Pharmacol*. 2001;134(4):845-852.
- 33. Stern CAJ, da Silva TR, Raymundi AM, et al. Cannabidiol disrupts the consolidation of specific and generalized fear memories via dorsal hippocampus CB1 and CB2 receptors. *Neuropharmacology*. 2017;125:220-230.

- Laprairie RB, Bagher AM, Kelly MEM, Denovan-Wright EM. Cannabidiol is a negative allosteric modulator of the cannabinoid CB1 receptor. *Br J Pharmacol.* 2015;172(20):4790-4805.
- Demuth DG, Molleman A. Cannabinoid signalling. *Life Sci.* 2006;78(6):549-563.
- 36. Johnson JR, Burnell-Nugent M, Lossignol D, Ganae-Motan ED, Potts R, Fallon MT. Multicenter, double-blind, randomized, placebo-controlled, parallel-group study of the efficacy, safety, and tolerability of THC:CBD extract and THC extract in patients with intractable cancer-related pain. *J Pain Symptom Manage*. 2010;39(2):167-179.
- Johnson JR, Lossignol D, Burnell-Nugent M, Fallon MT. An open-label extension study to investigate the long-term safety and tolerability of THC/CBD oromucosal spray and oromucosal THC spray in patients with terminal cancer-related pain refractory to strong opioid analgesics. *J Pain Symptom Manage*. 2013;46(2):207-218.
- Lichtman AH, Lux EA, McQuade R, et al. Results of a double-blind, randomized, placebo-controlled study of nabiximols oromucosal spray as an adjunctive therapy in advanced cancer patients with chronic uncontrolled pain. *J Pain Symptom Manage*. 2018;55(2):179-188.e1.
- 39. Canadian Agency for Drugs and Technologies in Health. Cannabinoid Buccal Spray for Chronic Non-Cancer or Neuropathic Pain: A Review of Clinical Effectiveness, Safety, and Guidelines. Ottawa, ON: Canadian Agency for Drugs and Technologies in Health; 2016.
- 40. Fitzcharles MA, Baerwald C, Ablin J, Häuser W. Efficacy, tolerability and safety of cannabinoids in chronic pain associated with rheumatic diseases (fibromyalgia syndrome, back pain, osteoarthritis, rheumatoid arthritis): A systematic review of randomized controlled trials. *Schmerz*. 2016;30(1):47-61.
- Johal H, Devji T, Chang Y, Simone J, Vannabouathong C, Bhandari M. Cannabinoids in chronic non-cancer pain: a systematic review and meta-analysis. *Clin Med Insights Arthritis Musculoskelet Disord*. 2020;13. https://doi.org/10.1177/ 1179544120906461.
- Erridge S, Salazar O, Kawka M, et al. An initial analysis of the UK Medical Cannabis Registry: Outcomes analysis of first 129 patients. *Neuropsychopharmacol Reports*. 2021;(April):1-9.
- von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies. 2015;147(8):573-578.
- 44. Quan H, Li B, Couris CM, et al. Updating and validating the Charlson Comorbidity Index and score for risk adjustment in hospital discharge abstracts using data from 6 countries. *Am J Epidemiol.* 2011;173(6):676-682.
- Brusselaers N, Lagergren J. The Charlson Comorbidity Index in registry-based research. *Methods Inf Med.* 2017;56(5):401-406.
- Herdman M, Gudex C, Lloyd A, et al. Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res.* 2011;20(10):1727-1736.
- Löwe B, Decker O, Müller S, et al. Validation and standardization of the generalized anxiety disorder screener (GAD-7) in the general population. *Med Care*. 2008;46(3):266-274.
- Snyder E, Cai B, DeMuro C, Morrison MF, Ball W. A new single-item sleep quality scale: results of psychometric evaluation in patients with chronic primary insomnia and depression. *J Clin Sleep Med.* 2018;14(11):1849-1857.
- 49. Pelayo-Alvarez M, Perez-Hoyos S, Agra-Varela Y. Reliability and concurrent validity of the palliative outcome scale, the

Rotterdam symptom checklist, and the brief pain inventory. J Palliat Med. 2013;16(8):867-874.

- Melzack R. The McGill Pain Questionnaire: major properties and scoring methods. *Pain*. 1975;1(3):277-299.
- Wolfe F, Clauw DJ, Fitzcharles MA, et al. 2016 revisions to the 2010/2011 fibromyalgia diagnostic criteria. *Semin Arthritis Rheum*. 2016;46(3):319-329.
- Van Hout B, Janssen MF, Feng YS, et al. Interim scoring for the EQ-5D-5L: mapping the EQ-5D-5L to EQ-5D-3L value sets. *Value Heal*. 2012;15(5):708-715.
- 53. National Institute for Health and Care Excellence (NICE) Position statement on use of the EQ-5D-5L value set for England. https://www.nice.org.uk/about/what-we-do/our-pro grammes/nice-guidance/technology-appraisal-guidance/ eq-5d-5l. Accessed July 10, 2021.
- Trotti A, Colevas AD, Setser A, et al. CTCAE v3.0: Development ment of a comprehensive grading system for the adverse effects of cancer treatment. *Semin Radiat Oncol.* 2003;13(3):176-181.
- 55. Ueberall MA, Essner U, Mueller-Schwefe GHH. Effectiveness and tolerability of THC:CBD oromucosal spray as add-on measure in patients with severe chronic pain: analysis of 12week open-label real-world data provided by the German pain e-registry. J Pain Res. 2019;12:1577-1604.
- Babson KA, Sottile J, Morabito D. Cannabis, cannabinoids, and sleep: a review of the literature. *Curr Psychiatry Rep.* 2017;19(4):23. https://doi.org/10.1007/s11920-017-0775-9.
- Whiting PF, Wolff RF, Deshpande S, et al. Cannabinoids for medical use: a systematic review and meta-analysis. JAMA. 2015;313(24):2456-2473.
- Safakish R, Ko G, Salimpour V, et al. Medical cannabis for the management of pain and quality of life in chronic pain patients: a prospective observational study. *Pain Med (United States)*. 2020;21(11):3073-3086.
- Goldenberg M, Reid MW, IsHak WW, Danovitch I. The impact of cannabis and cannabinoids for medical conditions on healthrelated quality of life: a systematic review and meta-analysis. *Drug Alcohol Depend*. 2017;174:80-90.
- Mücke M, Phillips T, Radbruch L, Petzke F, Häuser W. Cannabis-based medicines for chronic neuropathic pain in adults. *Cochrane Database Syst Rev.* 2018;2018(3).
- 61. Haroutounian S, Hohmann AG, Krane E, et al. Systematic review and meta-analysis of cannabinoids, cannabis-based medicines, and endocannabinoid system modulators tested for antinociceptive effects in animal models of injury-related or pathological persistent pain. *Pain.* 2021;162(suppl 1):S26-S44. https://doi.org/10.1097/j.pain.00000000002269.
- 62. Etges T, Karolia K, Grint T, et al. An observational postmarketing safety registry of patients in the UK, Germany, and Switzerland who have been prescribed Sativex® (THC: CBD, nabiximols) oromucosal spray. *Ther Clin Risk Manag.* 2016;12:1667-1675.
- Oreja Guevara C. Observational safety study of THC: CBD oromucosal spray (Sativex) in multiple sclerosis patients with spasticity. *Clin Exp Pharmacol.* 2015;5(184):2161-1459.

Supplemental Information

Additional supplemental information can be found by clicking the Supplements link in the PDF toolbar or the Supplemental Information section at the end of web-based version of this article.